



Recombinant Mouse IL-6

Catalog #	EPT096
Expression Host	E.coli
DESCRIPTION	Recombinant Mouse Interleukin-6 is produced by our E.coli expression system and the target gene encoding Phe25-Thr211 is expressed.
Accession	P08505
Synonyms	Interleukin-6; IL-6; B-Cell Hybridoma Growth Factor; Interleukin HP-1; Il6; Il-6
Mol Mass	21.8 KDa
AP Mol Mass	22 KDa, reducing conditions
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Endotoxin	Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
FORMULATION	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
RECONSTITUTION	Always centrifuge tubes before opening. Do not mix by vortex or pipetting.





It is not recommended to reconstitute to a concentration less than 100µg/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

STORAGE

Lyophilized protein should be stored at $< -20^{\circ}\text{C}$, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at $4-7^{\circ}\text{C}$ for 2-7 days.

Aliquots of reconstituted samples are stable at $< -20^{\circ}\text{C}$ for 3 months.

BACKGROUND

Interleukin-6 (IL-6) is a pro-inflammatory cytokine that also has an important role in immunity. Mouse IL-6 appears to be directly involved in the responses that occur after infection and injury and may prove to be as important as IL-1 in regulating the acute phase response. Mouse IL-6 is reported to be produced by fibroblasts, activated T cells, activated monocytes or macrophages, and endothelial cells. It acts upon a






ELK Biotechnology

variety of cells, including fibroblasts, myeloid progenitor cells, T cells, B cells and hepatocytes. IL-6 has a wide variety of biological functions: it plays an essential role in the final differentiation of B-cells into Ig-secreting cells, it induces myeloma and plasmacytoma growth, nerve cells differentiation in hepatocytes, and acute phase reactants.

SDS-PAGE



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei, P.R.C