



Recombinant Human TREM-1 (C-6His)

Catalog #	EPT113
Expression Host	Human Cells
DESCRIPTION	Recombinant Human Triggering Receptor Expressed on Myeloid Cells 1 is produced by our Mammalian expression system and the target gene encoding Ala21-Arg200 is expressed with a 6His tag at the C-terminus.
Accession	Q9NP99
Synonyms	Triggering Receptor Expressed on Myeloid Cells 1; TREM-1; Triggering Receptor Expressed on Monocytes 1; CD354; TREM1
Mol Mass	21.3 KDa
AP Mol Mass	32-40 KDa, reducing conditions
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Endotoxin	Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
FORMULATION	Lyophilized from a 0.2 μm filtered solution of 20mM





PB, 150mM NaCl, pH 7.2.

RECONSTITUTION

Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

It is not recommended to reconstitute to a concentration less than 100 μ g/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

STORAGE

Lyophilized protein should be stored at $< -20^{\circ}\text{C}$, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at $4-7^{\circ}\text{C}$ for 2-7 days.

Aliquots of reconstituted samples are stable at $< -20^{\circ}\text{C}$ for 3 months.

BACKGROUND

Triggering Receptor Expressed on Myeloid Cells 1 (TREM-1) is a transmembrane protein with a single Ig-like domain. TREM-1 associates with the adapter protein, DAP12, to deliver an activating signal. TREM-1 is expressed on blood neutrophils and monocytes, and





the expression is up-regulated by bacterial LPS. TREM-1 is expressed at high levels on neutrophils of patients with microbial sepsis and in mice with a TREM-1/Fc fusion protein protected mice against LPS-induced shock. Human TREM-1 shares 42% sequence homology with mouse TREM-1.

SDS-PAGE

